

## What happens if the glue ear is not treated?

Doctors do not really know if any damage occurs to the ear or hearing if the glue ear is not treated. We usually advise treating the problem if it does not clear up on its own to avoid the risk of long-term damage to the ear and hearing or problems in later life with language skills.

## Is glue ear common in adults?

Glue ear is uncommon in adults. It can follow on from a bad head cold, flu or other viral infection of the ear, nose or sinuses. Rarely, it can be caused by a serious blockage of the tube that goes from the back of the nose to the ear. (The Eustachian tube). Adults with glue ear should be seen by a ENT specialist as soon as possible.

*If you have any problems or questions, please contact:*

*Please insert local department routine and emergency contact details here*

**Disclaimer:** *This publication is designed for the information of patients. Whilst every effort has been made to ensure accuracy, the information contained may not be comprehensive and patients should not act upon it without seeking professional advice.*

*Last updated: November 2012 - Review due: November 2015*

# Glue Ear (OME)

**By Peter Robb and Haytham Kubba**

ENT UK is the professional Association for British Ear, Nose and Throat Surgeons and related professionals. This leaflet provides some background information about Glue Ear. It may be helpful in the discussions you have with your GP or specialist when deciding on possible treatment. This information leaflet is to support and not to substitute the discussion between you and your doctor. Before you give your consent to the treatment, you should raise any concerns with your GP or specialist.



**ENT UK**  
The Royal College of Surgeons of England  
35-43 Lincoln's Inn Fields  
London WC2A 3PE



ENT UK trading as British Academic Conference in Otolaryngology (BACO) and British Association of Otorhinolaryngology - Head and Neck Surgery (BAO-HNS)  
Registered as a Company limited by Guarantee in England and Wales under Company No 06452601  
Registered with the Charity Commission in England and Wales under Charity No 1125524

## Introduction

Glue ear is common. Up to eight in every ten children (80%) will have a short episode of glue ear before they start primary school. The medical name for glue ear is “otitis media with effusion”.

Doctors are not sure about all the causes of glue ear. Sometimes it follows after an ear infection, but many children with glue ear have never had an ear infection. The adenoid in the back of the nose may become infected with coughs and colds and the bacteria spread into the ear causing inflammation. The fluid (or “glue”) probably forms in the ear as a result of this inflammation.

Often, the hearing loss from glue ear is not enough to be noticed by the parents. Often, it is the child minder or nursery teacher who notices that the child cannot hear that well in a group situation. Sometimes, in a younger child, the hearing does not seem to be a problem, but the child’s speech and language development is slower than his or her friends of the same age.

Some children complain of earache because of the fluid in their ears. Some children have balance problems or poor attention as a result of glue ear. Sometimes, the only problem reported by parents or carers is behavioural problems. This is probably due to frustration on the part of the child who cannot hear properly.

If you have concerns about your child’s hearing or speech and language development, you should ask your health visitor or GP to refer your child for a hearing test. For children under the age of four years, this will probably be at a community hearing clinic. For older children, they will probably be referred to an ENT (ear, nose and throat) clinic in a hospital to see a specialist and have a hearing test.

For most children, the glue ear will get better with no treatment. You will probably be asked to come back for a second hearing test three or four months after the first test. Many children will get better over this time.

Those children who do still have problems after this period of what doctors call ‘watchful waiting’ or ‘active monitoring’ will probably be recommended surgical treatment. This may be grommet surgery or adenoidectomy and grommet surgery.

## Frequently asked questions

### Is there any treatment that will help the glue ear clear away more quickly?

The evidence is that neither medical treatments such as antibiotics or antihistamines nor alternative treatments such as cranial osteopathy are any better than waiting for a period of three months to see if the glue ear clears on its own.

### If my child has glue ear, are there any alternatives to grommets?

Hearing aids will help the hearing and give more time for the glue ear to clear. You can discuss this with your specialist.

### Should some children with glue ear have hearing aids as a first treatment?

For some children, glue ear can be a problem for much longer than others. In children with Down’s syndrome or cleft palate, hearing aids should be discussed with your specialist as a first treatment for glue ear.

### Is there anywhere else I can get good information about glue ear?

In 2008, the National Institute for Health and Clinical Excellence (NICE) published a guideline about the treatment of glue ear in children. [[www.nice.org.uk](http://www.nice.org.uk)].